

## Proceedings of the International Astronomical Union

Date of delivery: 5 May 2016

Journal and vol/article ref: IAU 1600053

Number of pages (not including this page): 3

This proof is sent to you on behalf of Cambridge University Press. Please check the proofs carefully. Make any corrections necessary on a hardcopy and answer queries on each page of the proofs

Please return the **marked proof** within 5 days of receipt to:

Managing editor of this symposium

**Authors are strongly advised to read these proofs thoroughly because any errors missed may appear in the final published paper. This will be your ONLY chance to correct your proof. Once published, either online or in print, no further changes can be made.**

To avoid delay from overseas, please send the proof by airmail or courier.

If you have **no corrections** to make, please email **managing editor** to save having to return your paper proof. If corrections are light, you can also send them by email, quoting both page and line number.

- The proof is sent to you for correction of typographical errors only. Revision of the substance of the text is not permitted, unless discussed with the editor of the journal. Only **one** set of corrections are permitted.
- Please answer carefully any author queries.
- Corrections which do NOT follow journal style will not be accepted.
- A new copy of a figure must be provided if correction of anything other than a typographical error introduced by the typesetter is required.

**If you do not send any corrections to the editor within 5 days, we will assume your proof is acceptable.**

- If you have problems with the file please contact

**lwebb@cambridge.org**

Please note that this pdf is for proof checking purposes only. It should not be distributed to third parties and may not represent the final published version.

**Important:** you must return any forms included with your proof. We cannot publish your article if you have not returned your signed copyright form.

NOTE - for further information about **Journals Production** please consult our **FAQs** at [http://journals.cambridge.org/production\\_faqs](http://journals.cambridge.org/production_faqs)

---

**Author queries:**

---

**Typesetter queries:**

---

**Non-printed material:**

# The Lyman- $\alpha$ Solar Telescope (LST) for the ASO-S Mission

Hui Li

Purple Mountain Observatory, Chinese Academy of Sciences, 2 West Beijing Road, Nanjing  
210008, China

email: nj.lihui@pmo.ac.cn

**Abstract.** The Lyman- $\alpha$  ( $Ly\alpha$ ) Solar Telescope (LST) is one of the payloads for the proposed Space-Borne Advanced Solar Observatory (ASO-S). LST consists of a Solar Disk Imager (SDI) with a field-of-view (FOV) of  $1.2 R_{\odot}$  ( $R_{\odot}$  = solar radius), a Solar Corona Imager (SCI) with an FOV of  $1.1 - 2.5 R_{\odot}$ , and a full-disk White-light Solar Telescope (WST) with the same FOV as the SDI, which also serves as the guiding telescope. The SCI is designed to work in the  $Ly\alpha$  (121.6 nm) waveband and white-light (for polarization brightness observation), while the SDI will work in the  $Ly\alpha$  waveband only. The WST works in both visible (for guide) and ultraviolet (for science) broadband. The LST will observe the Sun from disk-center up to  $2.5 R_{\odot}$  for both solar flares and coronal mass ejections with high tempo-spatial resolution

**Keywords.** solar flare, coronal mass ejection, coronagraph

---

## 1. Introduction

Solar flares and coronal mass ejections (CMEs) are the most energetic eruptions on the Sun and the primary drivers of disturbed (disastrous) space weather (Schwenn 2006). During flaring processes, particles (electrons, protons, neutrons, etc.) are accelerated to very high energy (MeV to GeV). After the initiation of CMEs, they are accelerated in a short time to high speed (up to more than 2000km/s) and may move towards the Earth with vast mass and numerous particles. When high energy particles and vast mass come close to the Earth, they will have great impact on the Earth's environment and subsequently on our daily life and the property in space.

The neutral hydrogen  $Ly\alpha$  line is the brightest EUV coronal line and very powerful in diagnosing structures of the solar atmosphere. As many proposed instruments, such as the Extreme Ultraviolet Imager (EUI) and the Multi Element Telescope for Imaging and Spectroscopy (METIS) on Solar Orbiter, the Chromospheric Lyman-Alpha SpectroPolarimeter (CLASP) and the Lyman Alpha Spicule Observatory (LASSO), LST will observe the Sun in the  $Ly\alpha$  line.

## 2. Scientific Objectives

The mission of the LST payload is to continuously monitor the Sun's activities and the source regions of space weather drivers, and to obtain high tempo-spatial resolution images of the Sun from the disk-center to  $2.5R_{\odot}$  in the  $Ly\alpha$  waveband and white-light (for polarization brightness). Its scientific objectives are delineated as follows.

- **To explore the relationship between CMEs and eruptive prominences and solar flares.** This relationship has been long in debate and so far inconclusive. Statistical studies show that there exists clear association between flares and CMEs (e.g., Wang & Zhang 2007), and the association rate increases with the energy of events (Yashiro & Gopalswamy 2009). LST will image the solar disk and the inner corona




Figure 1. Optical configuration of the SCI (upper left) and SDI (lower left), and the specifications of the LST instrument.

simultaneously in the same waveband and therefore is quite suitable for conducting such a study.

- **To study the triggering mechanisms, physical parameters and evolutions of solar flares and CMEs.** Even though past studies have narrowed down the possible CME initiation models to two main classes: those in which the pre-eruptive current-carrying magnetic field structures are sheared arcades (e.g., Antiochos *et al.* 1999) and those in which they are flux ropes (e.g., Amari *et al.* 2014), we are still far way from the conclusion. It is still a challenges to understand the triggering and energy accumulating mechanisms of these events in the low solar corona.

- **To investigate the acceleration mechanisms of CMEs in the inner corona and their kinetic behaviors.** Once started, CMEs often show quick expansion and acceleration, which is linked to both the rising phase of the soft X-ray flare (Zhang *et al.* 2001) and the particle acceleration (Berkebile-Stoiser *et al.* 2012). LST will provide the full-coverage data of the solar disk and the inner corona to address these scientific questions.

- **To determine the source regions of CMEs and their physical property.** Both solar flares and CME take place in the very low layer of the solar atmosphere. The physical conditions of the source regions of flares and CMEs on the disk determine, in some degree, their behavior and characteristics. LST together with the Full-disk vector Magnetograph (FMG) payload on the ASO-S will easily locate the source regions and the observation data are quite suitable for their property study.

### 3. Instrumentation

I give here the composition, up-to-date designs, specifications and key technical issues of the LST instrument in the following subsections, respectively.

#### 3.1. Composition & Current Designs

The LST is composed of a Solar Disk Imager (SDI), a Solar Corona Imager (SCI) and a full-disk White-light Solar Telescope (WST). This combination can well address the above-mentioned scientific objectives. Please refer to Figure 1 (right) for details of the instrument specifications.

71 The SCI is an inner-occulted Lyot-type reflective coronagraph. Its optical design is  
72 shown in Figure 1 (upper left). For SDI, two mirrors are used to image the Sun on the  
73 focal plan (Figure 1, lower left). A partial transmission filter is used to split the incident  
74 light of the WST and subsequently feed the guiding and polarization brightness channels.  
75 We do not show here the optical layout of the WST to save space.

### 76 3.2. Key technical issues

77 The most important two issues for the LST development (especially for the SCI) are  
78 stray-light control and image stability. Conjugate design, baffles, light-trap and super-  
79 polished mirror are used to realize the required stray-light level. Of all the stray-light  
80 sources, scattering from the main mirror is most significant stray-light source. To achieve  
81 the required stray-light, the required RMS surface roughness for the main is about 0.1 -  
82 0.2 nm, which is of challenge but possible. Cleanroom needs also to be used to minimize  
83 the contributions of dusts and particles during the developing processes.

84 The WST also serves as the guiding telescope and produces a white-light image of  
85 the solar disk on the focal plan, where four Photo Multiplier Tubes (PMTs) are placed  
86 to detect the solar limb and provide the guiding signal via some processing unit. After  
87 processing and calculation, the signal is used to drive the PZTs behind the main mirror  
88 to stabilize the image. Experiments demonstrate that this can well achieve the designed  
89 stability.

90 Other key issues include the fast readout of large format CCD camera with low noise,  
91 low temperature maintenance of the CCD chip, and accurate control of the PZTs, etc.

## 92 4. Summary

93 The ASO-S and its proposed payloads including LST are now under phase B study  
94 and is proposed to be launched around the next solar maximum. According to the design  
95 and outlined scientific objectives, LST will open a new window ( $\text{Ly}\alpha$ ) to observe the  
96 Sun continuously from disk center up to  $2.5R_{\odot}$  without gap with high tempo-spatial  
97 resolution, and contribute to the study of solar flares, CME initiation, evolution and  
98 kinetics, of the relation between flares, prominence eruptions and CMEs, and of the  
99 interface region between the corona and the chromosphere, etc.

## 100 Acknowledgements

101 The LST derives heritage from the Lyman- $\alpha$  Orbital Telescope (LYOT) for the once  
102 proposed French-Chinese SMall Explorer for Solar Eruptions (SMESE) (Vial *et al.* 2007)  
103 mission (phase A/A+ study finished). The LST related works are partially supported by  
104 NSFC via grant 11273065 and 11427803, and the Strategic Priority Research Program  
105 of Space Science of CAS via grant XDA04061003 and XDA04076101.

## 106 References

- 107 Amari, T. & Canou, A., Aly J.-J. 2014, *Nature*, 514, 465  
108 Antiochos, S. K. *et al.* 1999, *ApJ*, 510, 485  
109 Berkebile-Stoiser, S. *et al.* 2012, *ApJ*, 753, 88  
110 Schwenn, R. 2006, *Living Rev. Solar Phys.*, 3, 2  
111 Vial, J.-C., Auchere, F., & Chang, J. *et al.* 2007, *Adv. space Res.*, 40, 1787  
112 Wang, J. X. & Zhang, J. 2007, *Adv. space Res.*, 40, 1770  
113 Yashiro, S. & Gopalswamy, N. 2009, *IAUS*, 257, 233  
114 Zhang, J. & Dere, K, Howard, R. *et al.* 2001, *ApJ*, 559, 452