

Proceedings of the International Astronomical Union

Date of delivery: 5 May 2016

Journal and vol/article ref: IAU 1600209

Number of pages (not including this page): 3

This proof is sent to you on behalf of Cambridge University Press. Please check the proofs carefully. Make any corrections necessary on a hardcopy and answer queries on each page of the proofs

Please return the **marked proof** within **5** days of receipt to:

Managing editor of this symposium

Authors are strongly advised to read these proofs thoroughly because any errors missed may appear in the final published paper. This will be your ONLY chance to correct your proof. Once published, either online or in print, no further changes can be made.

To avoid delay from overseas, please send the proof by airmail or courier.

If you have **no corrections** to make, please email **managing editor** to save having to return your paper proof. If corrections are light, you can also send them by email, quoting both page and line number.

- The proof is sent to you for correction of typographical errors only. Revision of the substance of the text is not permitted, unless discussed with the editor of the journal. Only **one** set of corrections are permitted.
- Please answer carefully any author queries.
- Corrections which do NOT follow journal style will not be accepted.
- A new copy of a figure must be provided if correction of anything other than a typographical error introduced by the typesetter is required.

If you do not send any corrections to the editor within 5 days, we will assume your proof is acceptable.

- If you have problems with the file please contact

lwebb@cambridge.org

Please note that this pdf is for proof checking purposes only. It should not be distributed to third parties and may not represent the final published version.

Important: you must return any forms included with your proof. We cannot publish your article if you have not returned your signed copyright form.

NOTE - for further information about **Journals Production** please consult our **FAQs** at http://journals.cambridge.org/production_faqs

Author queries:

Typesetter queries:

Non-printed material:

Multitemperature analysis of solar flare observed on 2003 March 29

Anna Kepa¹, Barbara Sylwester¹, Janusz Sylwester¹, Marek
Siarkowski¹, Tomasz Mrozek^{1,2} and Magdalena Gryciuk^{1,2}

¹Space Research Centre, Polish Academy of Sciences, Solar Physics Division, Wrocław, Poland
email: ak@cbk.pan.wroc.pl

²Astronomical Institute, University of Wrocław, Wrocław, Poland

Abstract. We present results of multitemperature analysis of *GOES* C7.2 class flare SOL2003-03-29T10:15. This event occurred close to the centre of the solar disk and had two maxima in soft X-rays. We have performed analysis of physical parameters characterizing evolution of conditions in the flaring plasma. The temperature diagnostics have been carried out using the differential emission measure (DEM) approach based on the soft X-ray spectra collected by RESIK Bragg spectrometer. Analysis of data obtained by *RHESSI* provided opportunity to estimate the volume and thus calculating the density and thermal energy content of hot flaring plasma.

Keywords. Solar flares, spectra, differential emission measure

1. Introduction

During the RESIK experiment (Sylwester *et al.*, 2005) onboard the *CORONAS-F* satellite several thousand of X-ray spectra in the wavelength range from 3.3 to 6.5 Å were obtained. The measurements include spectra of solar flares, active regions and quiet corona. The analysis of selected lines intensities allows the study of physical conditions in solar plasma. In this paper we focus our attentions on determinations and analysis of differential emission measure (DEM $\equiv \varphi(T_e)$) distributions, which is defined as follows: $\varphi(T_e) = N_e^2 \frac{dV}{dT_e}$ (N_e - electron density, V - plasma volume, T_e - temperature). The DEM convolved with the theoretically determined emission function ($f(T_e)$) of each selected wavelength gives the observed flux in appropriate spectral line/band i .

$$F_i = A_i \int_0^\infty f_i(T_e) \varphi(T_e) dT_e \quad i = 1, 2, \dots, N \quad (1.1)$$

where: A_i represents the assumed abundance of an element contributing to the flux of a particular line or spectral interval and N is the number of spectral bands used. The reconstruction of the DEM from measured line fluxes is based on the inversion of the set of integral equations 1.1 for N selected lines bands. To obtain a reliable DEM a set of lines with emission functions widely distributed over the temperature interval is required. Unfortunately a direct inversion of the data does not produce a unique DEM solution and additional constraints are needed to achieve a stable solution of this ill-posed inverse problem. There are a number of existing algorithms for reconstructing DEMs from solar and stellar data (e.g. Sylwester *et al.*, 1980; Aschwanden *et al.*, 2015). In this contribution we present (for the first time) the differential emission measure distributions obtained based on RESIK spectra using the Adaptive Differential Evolution method.




Figure 1. Left: *GOES* 1 - 8 Å lightcurve (in red) with optimal fit of elementary flares (thin black lines). The thick black line represents the sum of individual overlapped flare profiles. The grey strips indicate time intervals of passages through a polar van Allen radiation belt when the RESIK high-voltages were turned off and no observations were made. The four numbers denote intervals over which RESIK spectra were integrated for DEM analysis. Right: Average RESIK spectrum (integrated over 24 min) for SOL2003-03-29T10:15 flare. The lines used in the analysis are marked (see Table 1 for the details). The dashed lines limit the four RESIK channels bands. Unfortunately, due to technical problems during this flare, channel 3 data are not reliable for the analysis.

Table 1. Spectral bands used to calculate the DEM distributions.

No.	Wavelength range [Å]	Main line	Channel
1	3.358 - 3.388	Ar XVII 3p	1
2	3.515 - 3.585	K XVIII (w)	1
3	3.715 - 3.754	Ar XVIII 2p	1
4	3.940 - 3.975	Ar XVII (w)	2
5	3.980 - 4.010	Ar XVII (z)	2
6	4.070 - 4.100	S XV 4p	2
7	5.000 - 5.075	S XV 2p (w)	4
8	5.075 - 5.150	S XV 2p (z)	4
9	5.180 - 5.250	Si XIV 3p	4
10	5.260 - 5.310	Si XIII 5p	4
11	5.350 - 5.430	Si XIII 4p	4
12	5.650 - 5.710	Si XIII 3p	4

2. Observations and analysis

We present results of analysis of C7.2 flare SOL2003-03-29T10:15 observed by RESIK. The flux from *GOES* shows that the flare is complex and consists of a number of successive individual flares. Using elementary soft X-ray flare temporal profile we selected 6 individual flares composing *GOES lightcurve* (Figure 1, left; Gryciuk *et al.*, 2015).

For four selected time intervals (corresponding to different flares) we calculated the mean spectrum and fluxes in $N = 12$ wavelength ranges (Table 1). To avoid the contribution of non-flaring plasma, the preflare X-ray fluxes have been subtracted.

DEM determinations were made using Adaptive Differential Evolution method (genetic algorithm). Differential evolution, introduced by Storn, & Price (1997) is simple but powerful evolutionary algorithm for global optimization. Starting from randomly chosen initial populations of different DEMs a new generation of DEMs is produced by crossover and mutations. Our population had 100 individual DEM distributions or 'chromosomes'. Each chromosome consists of 100 gens. They correspond to DEM values for 100 temperatures in the range from 2 to 30 MK. Process of breeding (and multiplication) of the whole population is controlled by assumed fitness criterion based on the values of observed to calculated fluxes. Based on the actual DEM distribution the predicted flux was calculated using the formula 1.1. For chlorine abundance we adopted




Figure 2. The DEM distributions obtained for individual intervals defined in Figure 1.

57 $A_{Cl} = 5.62 \times 10^{-7}$. For silicon, sulfur, argon, and potassium we used abundances calcu-
 58 lated using the multithermal assumption (AbuOpt method; Sylwester *et al.*, 2015). We
 59 assumed $A_{Si} = 2.618 \times 10^{-5}$, $A_S = 7.413 \times 10^{-6}$, $A_{Ar} = 3.083 \times 10^{-6}$, $A_K = 6.067 \times 10^{-7}$.
 60 For other elements we adopted abundances called as sun_coronal_ext.abund (available in
 61 the CHIANTI package). The process of evolution was stopped after 6000 generations,
 62 when the convergence became very slow. The minimum χ^2 values were in the range
 63 1.5 - 2.3. The process of evolution was repeated 10 times, each time starting from a new
 64 random population. Ten of the best DEM distributions obtained for individual intervals
 65 are presented in Figure 2.

66 For the maximum and decay phase of the first flare the hard X-ray measurements
 67 from *RHESSI* were available. *RHESSI* images were obtained with the PIXON algorithm
 68 in the energy range 6 - 8 keV. From this data we estimated volumes (spherical shape
 69 was assumed). The spatial dimensions combined with the total emission measure of the
 70 hotter component allowed to estimate the electron density and thermal energy content.

71 3. Conclusion

72 For the first time we present the DEM distributions calculated using the Adaptive
 73 Differential Evolution method. Our DEM distributions are two components, which is
 74 consistent with previous results obtained using Withbroe-Sylwester method. The cooler
 75 component corresponds to plasma from 3 MK to 10 MK, hotter conforms the temperature
 76 range 12 - 25 MK. For the first flare the average volume (estimated from *RHESSI* data) is
 77 $1.9 \times 10^{25} \text{ cm}^3$. This leads to the following values of electron density and thermal energy
 78 content for the hot component of the plasma: $2.9 \times 10^{11} \text{ cm}^{-3}$ as $4.11 \times 10^{28} \text{ erg}$.

79 Acknowledgements

80 This research has been supported by Polish National Science Centre grants No. DEC-
 81 2013/09/N/ST9/02209, 2011/01/M/ST9/05878, and 2011/01/M/ST9/06096.

82 References

- 83 Aschwanden, M. J., Boerner, P., Caspi, A., McTiernan, J. M., Ryan, D., & Warren, H. P. 2015,
 84 *Sol. Phys.* (in press)
 85 Gryciuk, M., Siarkowski, M., & Gburek, S. *et al.* 2015, in preparation
 86 Storn, R. & Price, K. 1997, *Journal of Global Optimization*, 11, 341
 87 Sylwester, B., Phillips, K. J. H., Sylwester, J., & Kepa, A. 2015, *ApJ*, 805, 49
 88 Sylwester, J. & Gaicki, I., Kordylewski Z. *et al.* 2005, *Sol. Phys.*, 226, 45
 89 Sylwester, J., Schrijver, J., & Mewe, R. 1980, *Sol. Phys.*, 67, 285